



(REPLACEMENT SPECIFICATION 10/812,709)

UNITED STATES PLANT PATENT APPLICATION

of

L. PERNILLE AND MOGENS N. OLESEN

for

COMPACT FLORIBUNDA
ROSE PLANT NAMED

'Poulac014'

SUMMARY OF THE INVENTION

BOTANICAL CLASSIFICATION

Rosa hybrida

VARIETY DENOMINATION

5 'Poulac014'

The present invention constitutes a new and distinct variety of garden rose plant which originated from a controlled crossing between the female seed parent plant, an
10 un-named seedling, and the male pollen parent plant, an un-named seedling. The two parents were crossed during the summer of 1995 and the resulting seeds were planted in a controlled environment in Fredensborg, Denmark. The new variety is named 'Poulac014'.

15 The new variety may be distinguished from its unnamed female seed parent, by the following combination of characteristics:

1. While the seed parent has yellow flowers,
'Poulac04' has apricot flowers.
- 20 2. While the seed parent has acute flower petal bases, 'Poulac014' has rounded to acute flower petal base shapes.

The new variety may be distinguished from its male pollen parent by the following combination of
25 characteristics:

1. 'Poulac014' has larger leaves than those of the pollen parent.
2. 'Poulac014' has larger flowers than the pollen parent.

5 The objective of the hybridization of this rose variety was to create a new and distinct variety for garden use with unique qualities, such as:

1. Uniform and abundant apricot flowers;
2. Vigorous, but compact growth when propagated
10 both as a budded rose and on its own roots;
3. Disease resistance.

 This combination of qualities is not present in previously available commercial cultivars of this type, known to the inventors, and distinguish 'Poulac014' from all
15 other varieties of which we are aware.

 As part of their rose development program, L. Pernille Olesen and Mogens N. Olesen germinated the seeds from the aforementioned hybridization during winter of 1995 and conducted evaluations on the resulting seedlings in a
20 controlled environment in Fredensborg, Denmark.

 'Poulac014' was selected in the spring 1996 by the inventors as a single plant from the progeny of the aforementioned hybridization.

 Asexual reproduction of 'Poulac014' by traditional
25 budding and rooted cuttings was first done by L. Pernille

and Mogens N. Olesen in their nursery in Fredensborg,
Denmark in July, 1996. This initial and other subsequent
asexual propagations conducted in controlled environments
have demonstrated that the characteristics of 'Poulac014'
5 are true to type and are transmitted from one generation to
the next.

BRIEF DESCRIPTION OF THE DRAWING

10 The accompanying color illustration shows as true as is
reasonably possible to obtain in color photographs of this
type, the typical characteristics of the buds, flowers,
leaves, and stems, of 'Poulac014'. Specifically illustrated
in the SHEET 1

15 Fig 1.1; Open flowers, above view and side view;
Fig 1.2; Flower bud closed, flower bud as sepals
unfold, and partially open;
Fig 1.3; Flower petals, detached;
Fig 1.4; Sepals, receptacle, and peduncle;
20 Fig 1.5; Juvenile leaf exhibiting anthocyanin;
Fig 1.6; Mature Leaf;
Fig 1.7; Bare stems, exhibiting thorns.

25

DETAILED DESCRIPTION OF THE VARIETY

The following is a description of 'Poulac014', as observed in its growth in a field nursery in Jackson County, Oregon. Observed plants are 3 years of age. Plants were grown on *Rosa multiflora* understock. Color references are made using the Royal Horticultural Society (London, England) Colour Chart, 1995, except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety 'Poulymp', a rose variety from the same inventors described and illustrated in U.S. Plant Patent Application No. 09/607,327 dated 30 June, 2000, are compared to 'Poulac014' in Chart 1.

CHART 1

	'Poulac014'	'Poulymp'
Flower Diameter	60 to 65 mm	60 to 70 mm
Petalage	30 to 35 petals	25 to 30 petals
Flower Color after opening: Upper surface of outermost petals.	Orange Group 25B at marginal to middle zone, becoming Yellow-Orange Group 14A at basal zone.	Yellow-Orange Group 21 C

FLOWER AND FLOWER BUD

Blooming habit: Continuous.

5 Flower bud:

Size: Upon opening, 28 mm in length from
base of receptacle to end of bud.
Bud diameter is 17 mm on average.

Bud form: Pointed ovoid with broad base.

10 Bud color: As sepals unfold, petals are
Orange Group 25D to Yellow Group
12C.

Sepals:

Upper Surface:

15 Color: Yellow-Green Group 144A
to 144B.

Surface: Moderately pubescent.

Lower Surface:

20 Color: Yellow-Green Group
144A.

Sepal Shape:

Sepal apex is cirrhose. Base is
flat at union with receptacle.

Sepal Margin:

25 Margins have strong foliaceous

appendages on three of the five
sepals.

Size: 28 mm (l) x 8 mm (w).

Receptacle:

5 Surface Texture:

Smooth to Glabrous.

Shape: Funnel shaped.

Size: 4 mm (h) x 8 mm (w).

Color: Yellow-Green Group 144B.

10 Pedicel:

Surface: Smooth. Few to medium
quantity of stipitate glands
towards base of peduncle.

Length: 25 to 30 mm.

15 Diameter: 3 mm on average.

Color: Yellow-Green Group 145B
Anthocyanic pigments the
color of Greyed-Red Group
180C observed.

20 Strength: Strong.

Borne: In clusters of 1 to 4 flower
buds per stem.

Flower bloom:

25 Fragrance: Moderate floral scent.

5	<p>Duration: The blooms have a duration on the plant of approximately 10 days. Petals fall cleanly away from plant after flowers have matured completely.</p>
10	<p>Size: Flower diameter is 60 to 65 mm when open. Flower depth is 30 to 35 mm.</p>
	<p>Form:</p> <p style="padding-left: 40px;">General shape is a deep cup.</p>
15	<p>Shape of flower when viewed from the side:</p> <p>Upon opening, upper part: Flat.</p> <p>Upon opening, lower part: Flat.</p> <p>Open flower, upper part: Flat.</p> <p>Open flower, lower part: Concave.</p>
20	<p>Petalage: Average range is 30 to 35 petals under normal conditions with 7 petaloids.</p>
	<p>Color:</p> <p>Upon opening, petals:</p> <p style="padding-left: 40px;">Outermost petals:</p>
25	<p style="padding-left: 40px;">Outer side: Orange-Red Group 34A at</p>

marginal to middle zones,
becoming Yellow Group 12A to
12B at basal zone.

5 Inner Side: Orange Group 25B at marginal
to middle zone, becoming
Yellow-Orange Group 14A at
basal zone.

Innermost petals:

10 Outer side: Orange-Red Group 30A to
Orange-Red Group 34B at
marginal to middle zones,
becoming Yellow-Orange Group
14A at basal zone.

15 Inner Side: Orange Group 26A, becoming
Yellow-Orange Group 14A at
basal zone.

20 **General Tonality:** On open flower Orange Group 25B
with intonations of Orange-Red
Group 33B to 33C. No change in the
general tonality at the end of the
10th day. Afterwards, general
tonality is Yellow-Orange Group
23C to 23D.

25

Petals:

Petal Reflex: Outer petals are somewhat
reflexed.

5 Margin: Entire and uniform with an
occasional cleft. Medium
undulations of margin observed.

Shape: Apex: Round.
Base: Round to acute.

Size: 35 mm (l) x 32 mm (w).

10 Texture: Smooth.

Thickness: Thick.

Arrangement: Not Formal.

Petaloids:

15 Quantity: 5 to 9.

Color:

Upper Surface: Yellow-Orange Group 23A to
Yellow Group 12A.

Lower Surface: Orange-Red Group 34C to
20 Yellow Group 12A.

Size: 20 mm (l) x 10 mm (w).

Shape: Apex is round. Base is rounded to
acute.

25

Reproductive Organs:

Pistils:

Length: 8 mm.

Quantity: 39 (actual count).

5

Pollen:

None observed.

Anthers:

Size: 2 mm in length.

Color: Yellow Group 10A.

10

Quantity: 84 (actual count).

Filaments:

Color: Yellow Group 13A.

Length: 10 mm.

Stigmas: Inferior relative to the

15

length of the filaments and
the height of the anthers.

Color: Greyed-Yellow Group 162D.

Styles:

Color: Red-Purple Group 57A.

20

Length: 10 mm on average.

Hips: None Observed in the field nursery in
Jackson County Oregon.

PLANT

25

Plant growth: Compact, upright to bushy. When grown
as a budded field grown plant on Rosa
multiflora understock, the average
height of the plant is 60 cm and the
average width is 60 cm.

5

Stems:

Color:

Young wood: Yellow-Green Group 144C.

Older wood: Green Group 138B.

10

Surface Texture:

Young wood: Smooth.

Older wood: Smooth.

Thorns:

Incidence: 23 thorns per 10 cm of stem.

15

Size: Average length: 10 mm.

Mature color: Greyed-Yellow Group 162A.

Juvenile color: Yellow-Green Group 144C to Greyed-
Red Group 181A and 181B.

20

Some intonations of Greyed-Yellow
Group 162A observed.

Shape: Deeply concave.

Plant foliage: Normal number of leaflets on
normal leaves in middle of the
stem: 5 leaflets.

25

margins and lower side.

Color: Green Group 143A.

Petiole:

Length: 43 mm.

5 Diameter: 1 mm.

Above:

Color: Yellow-Green Group 144B.

Anthocyanin: Greyed-Red Group 181B.

Underneath:

10 Observations: Thorns, fragrant
stipitate glands and
light pubescence.

Rachis:

Length: 65 mm.

15 Above:

Color: Yellow-Green Group 144B

Underneath:

20 Observations: Thorns, fragrant
stipitate glands, and
light pubescence.

Leaflet:

Size: 65 mm (l) x 41 mm (w).

Edge: Shallow serrations.

25 Shape: Ovate. Apex is acute to

rounded. Base is rounded.

Texture: Smooth.

Arrangement: Odd pinnate.

Venation: Reticulate.

5 Glossiness: Glossy.

Thickness: Thick and leathery.

Disease resistance:

Above average resistance to mildew, rust, black spot,
10 and Botrytis under normal growing conditions in Jackson
County, Oregon.

Cold Hardiness:

The variety 'Poulac014' has been found to be cold
15 tolerant to USDA Cold Hardiness Zone 6.